# Thin points of a class of Markov processes with jumps

#### Xiaochuan YANG, Université Paris-Est

joint work with S. Seuret (Créteil, France)

CJPS





Houches, 21/4/2016

1/13

#### Basic problem

Question : regularity ?



#### Basic problem

- $X = \{X_t, t \in [0, 1]\}$  in  $\mathbb{R}^d$ .
- $\mu(A) = \int_0^1 \mathbf{1}_A(X_t) dt$  for all  $A \subset \mathbb{R}^d$ .

#### Question : regularity ?

- Absolutely continuity (local times when X is Markovian).
- ▶ Local dimensions, i.e. for  $x \in \text{supp}(\mu)$ , the positive real h such that

$$\mu(B(x,r)) \sim r^h.$$

- ► Always well defined, i.e.  $\lim_{r\to 0} \frac{\ln \mu(B(x,r))}{\ln r}$  exists?
- How does h depend on the value x? study a regularity exponent h(x).

## Examples

- B : Brownian motion in  $\mathbb{R}^d.$ 
  - ▶ d = 1: local times exist [Lévy].
  - ▶  $d \ge 2$ : local dimension is 2 for all  $x \in \text{supp}(\mu)$  [Perkins-Taylor].

## Examples

- B : Brownian motion in  $\mathbb{R}^d.$ 
  - ▶ d = 1 : local times exist [Lévy].
  - ▶  $d \ge 2$ : local dimension is 2 for all  $x \in \text{supp}(\mu)$  [Perkins-Taylor].
- $\sigma:\alpha\text{-stable subordinator, i.e.}$  increasing stable Lévy process in  $\mathbb{R}^+.$ 
  - ▶ Local dimension is  $\alpha$  for  $\mu$ -almost every  $x \in \text{supp}(\mu)$  [Hu-Taylor].

・ロン ・四 と ・ 日 ・ ・ 日 ・

• Exceptional points? Yes.

## Framework : multifractal analysis

**Goal** : distinguish different local behaviors of  $\mu$  by a description of the "size" of the set of points with given regularity.

Definition

The upper local dimension of  $\mu$  at x is defined by

$$\overline{h}(\mu, x) = \limsup_{r \to 0} \frac{\ln \mu(B(x, r))}{\ln r}.$$

4/13

One defines similarly the lower local dimension  $\underline{h}(\mu, x)$  and local dimension  $h(\mu, x)$  when the limit exists.

#### Definition

Define the iso-holder sets

$$\overline{E}(h) = \{ x \in \operatorname{supp}(\mu) : \overline{h}(\mu, x) = h \}.$$

The upper multifractal spectrum of  $\mu$  is the mapping

 $\overline{d}_{\mu}(\cdot): h \mapsto \dim_{\mathcal{H}} \overline{E}(h).$ 

One defines similarly  $\underline{d}_{\mu}(\cdot)$  and  $d_{\mu}(\cdot)$ .

"Recall" : Hausdorff dimension describes the size of "small" sets in a metric space, e.g. a triadic Cantor set in  $\mathbb{R}^1$ .

(ロ) (部) (目) (日) (日) (の)

#### Thin points for $\alpha$ -stable subordinator

**Recall** local dimension exists for typical points :  $h(\mu, x) = \alpha$  for  $\mu$ -almost every point in supp  $\mu$ , i.e.

$$\mu(B(x,r)) \sim r^{\alpha}.$$

However, there are "many" points with smaller than normal mass, i.e.

$$\mu(B(x,r)) \sim r^h \text{ with } h > \alpha.$$

These points are called thin points.

Theorem (Hu-Taylor)

A.s. the following holds

$$\overline{d}_{\mu}(h) = \begin{cases} \alpha(\frac{2\alpha}{h} - 1) & \text{ if } h \in [\alpha, 2\alpha], \\ -\infty & \text{ otherwise.} \end{cases}$$

Our process : stable-like jump diffusion

- ► **Goal** : describe thin points of jump diffusions (i.e. jumping SDE) by multifractal analysis.
- ► **Difference/Difficulty :** no more stationary increment, Markovian dynamic is space-dependent.

Definition (Bass)

 $The \ stable-like \ jump \ diffusion \ is \ a \ Markov \ processes \ with \ generator$ 

$$\mathcal{L}f(x) = \int_0^1 f(x+u) - f(x)\frac{du}{u^{1+\beta(x)}}$$

where  $\beta$  is a Lipschitz function taking value in  $[\varepsilon, 1 - \varepsilon]$ .

**Remark :** when  $\beta(\cdot) = \alpha \in (0, 1)$ , one recovers  $\alpha$ -stable subordinator (truncated large jumps).

The stable-like jump diffusion satisfies the jumping SDE

$$M_t = \int_0^t \int_0^1 z^{1/\beta(M_{s-1})} N(ds, dz).$$

where N(ds, dz) is a Poisson random measure with intensity  $\pi(dz) = dz/z^2$ .

**Remind** : dimension of the sets  $\overline{E}(h) = \{x \in \text{supp}(\mu) : \overline{h}(\mu, x) = h\}$ . Need a translation!

#### Preparation I : heuristic computation

As  $t \in \overline{E}(h)$ , necessarily

$$\mu(B(M_t, r_n)) \le r_n^{h-\varepsilon}, \text{ for } r_n \to 0.$$

 $\mu(\cdot)$  measures the time spent by M inside balls, last inequality means M can not move too slowly, precisely, infinitely often

$$|M_{t+2^{-n}} - M_t| \wedge |M_t - M_{t-2^{-n}}| \ge 2^{-n/((h-\varepsilon)\beta(M_t))}.$$

イロト イロト イヨト イヨト 三日

Need estimate for increments.

# Preparation II : a key estimate

#### Proposition

For all  $\delta > 1$ ,  $m \in \mathbb{N}^*$ , with probability larger than  $1 - e^{-m}$ , for  $|t - s| \sim 2^{-m}$ ,

$$\left| \int_{s}^{t} \int_{0}^{2^{-\frac{m}{\delta}}} z^{1/\beta(M_{u-})} N(du, dz) \right| \leq \log\left(\frac{1}{|s-t|}\right)^{2} |s-t|^{\frac{1}{\delta \cdot \beta_{s,t}^{m}}}$$
  
with  $\widehat{\beta}_{s,t}^{m} \approx \sup_{u \in [s,t]} \beta(M_{u-}).$ 

**Remark :** uniformly, small jumps accumulation has the same effect of a single jump.

So there are two "large" jumps beside  $t\in \overline{E}(h)$  for infinitely many time scales.

Highlight double jumps configuration in the Poisson point process gives an upper bound for  $\dim_{\mathcal{H}} \overline{E}(h)$ .

Lower bound is more involved.

## Multifractal spectrum

Theorem (16' Seuret and Y.)

A.s. the upper multifractal spectrum of  $\mu$  is



イロト イロト イヨト イヨト 三日

12/13

**Remark :** superposition of random curves.

Merci de votre attention !

